



Calcium

Calcium Plus is a superior Calcium formulation fortified with Plant Elicitors to enhance disease resistance.

Ca 17%

20L



9 418768 000124



200L



9 418768 000131



Calcium Plus Application Rates

Crop	Application rate (l/ha)	When to apply	Water rate (l/ha)
Apples & Pears	5 - 7	Up to 8 applications from bud burst to 30 days pre harvest at 10 to 12 day intervals. For apple varieties extremely prone to bitter pit, these application rates may be increased to 7 litres per hectare. Avoid spraying over flowering and for 2 weeks after.	500 - 1000
Curcubits & Capsicums	5	2 to 4 applications at 5 litres per hectare from 2 weeks after emergence at 10 day intervals.	200 - 500
Kiwifruit	5	3 to 4 applications of 5 litres per hectare from bud burst at 14 day intervals. Avoid spraying over flowering and for 2 weeks after.	500 - 1000
Brassicas, Silverbeet & Lettuce	5	3 applications of 5 litres per hectare at 10 day intervals commencing after 14 to 21 days from transplanting.	200 - 500
Citrus	7	3 to 4 applications of 7 litres per hectare at 14 day intervals commencing at bud burst. Avoid spraying over flowering and for 2 weeks after.	500 - 1000
Onions	7	3 to 4 applications of 7 litres per hectare at 14 day intervals commencing when there is sufficient leaf to accept the spray.	200 - 500
Carrots	5	2 to 3 applications of 5 litres per hectare at 14 day intervals commencing when there is sufficient leaf to accept the spray.	200 - 500
Peaches, Nectarines & Apricots	5 - 7	2 applications of 5 litres per hectare at 10 day intervals commencing after bud burst, followed by 2 to 3 applications of 7 litres per hectare at 14 to 21 day intervals to 30 days pre harvest. Avoid spraying over flowering and for 2 weeks after.	500 - 1000
Potatoes	5 - 7	2 applications of 5 litres per hectare at 10 day intervals from tuber initiation followed by one application of 7 litres per hectare.	200 - 500
Tomatoes	5 - 7	3 to 4 applications of 5 litres per hectare at 10 day intervals from fruitset on second truss.	200 - 500
Grapes	5 - 7	5 to 7 litres per hectare at 10 day intervals from fruitset to one month pre harvest.	500 - 1000

Mixing and Application:

- Shake the container before adding contents to spray tank. Agitate mix while filling and agitate regularly until spraying is completed. Calcium Plus liquid calcium has proven compatible with a wide range of other agrichemicals. Stability can be checked by adding a proportionate amount of Calcium Plus liquid calcium to a sample of previously mixed spray. When using the highest recommended application rates, also use the highest recommended water rates. Do not spray any crop in direct sunlight or at times of high temperature and rapid drying.
- Wear protective clothing and equipment when spraying.

- Ensure that spraying equipment is clean before use and that all equipment is washed thoroughly after use. Store in a safe place and don't allow prolonged exposure to high and low temperatures.

Conditions of sale:

The use of this product being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Omnia accept no responsibility whatsoever resulting from the use of this product.

Batch:

www.omnia.net.nz
 Freephone 0800 774 629
 PO Box 201041 Auckland Airport, 2150

