



# Hiyield

Hiyield Plus is NPK 5.4.11 with trace elements, seaweed extracts, surfactants and plant health activators to promote size, colour, yields and also to help reduce the effects of disease pressure in fruit and vegetable crops.

20L



9 418768 000087



200L



9 418768 000094

# Hiyield Plus Application Rates

Crop	Application rate (l/ha)	When to apply	Water rate (l/ha)
Apples, Nashi, Pears	3 - 7	Starting from fruit set until harvest, apply 3 l/ha on a 7-10 day basis or at 10 l/ha on a 4 weekly cycle.	500 - 1000
Avocados	5 - 7	Starting from six weeks after new fruit set until just prior to harvesting. Repeat applications on a 21 to 28 day cycle.	500 - 1000
Berryfruit	5 - 7	Apply at 21 day intervals from fruit set until two weeks prior to harvest.	500 - 1000
Citrus	5 - 7	Depending on varieties 3 to 4 weeks after fruit set until one month before harvest, repeat applications on a monthly basis.	500 - 1000
Grapes	5 - 7	Apply at 21 day intervals from fruit set until one month prior to harvest.	400 - 1000
Kiwifruit	5 - 7	Apply a total of 40 to 50 l/ha evenly spread over the growing season until just prior to harvest.	500 - 1000
Passionfruit	5 - 7	Apply every 7 to 14 days over the growing season from 2 weeks after final flowering until one month before harvest.	200 - 1000
Persimmons	5 - 7	Starting from fruit set until one month before harvest, apply 5 l/ha on a 7 to 10 day basis or at 7 l/ha on a weekly cycle until one month before harvest.	500 - 1000
Stone fruit	5 - 7	Apply a total of 30 to 40 l/ha evenly over the growing season from stone hardening to until one month before harvest.	500 - 1000
Soft fruit, Strawberries	5 - 7	Apply every 7 to 14 days from after the first fruit set through to the end of the season's harvest.	500 - 1500
Capsicums, Tomatoes	5 - 7	Apply a total of 40 to 50 l/ha evenly spread over the growing season from after the first fruit set through to the end of the season's harvest.	200 - 500
Cucurbits	5 - 7	Apply a total of 40 to 50 l/ha evenly spread over the growing season from after the first fruit set through to the end of the season's harvest.	200 - 500
Potatoes	5 - 7	Apply a total of 30 to 40 l/ha evenly spread over the growing season from after the first fruit set through to the end of the season's harvest.	200 - 500
Other		Call our free customer help line 0800 774 629	

## Mixing and Application:

- Shake the container before adding contents to spray tank
- Agitate mix while filling and agitate regularly until spraying is completed.
- Hiyield Plus has proven compatibility with a wide range of other agrichemicals.
- Stability can be checked by adding a proportionate amount of Hiyield Plus to a sample of previously mixed spray.

- When using the highest recommended application rates, also use the highest recommended water rates.
- Do not spray any crop in direct sunlight or at times of high temperature and rapid drying.
- Wear protective clothing and equipment when spraying.
- Ensure that spraying equipment is clean before use and that all equipment is washed thoroughly after use.
- Store in a safe place and do not allow prolonged exposure to high and low temperatures.

## Conditions of sale:

The use of this product being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Omnia accept no responsibility whatsoever resulting from the use of this product.

## Batch:

www.omnia.net.nz  
 Freephone 0800 774 629  
 PO Box 201041 Auckland Airport, 2150

