

HARMFUL

KEEP OUT OF REACH OF CHILDREN



A fungicide/bactericide for the control of leaf spot in olives, *Sclerotinia* fruit rot in kiwifruit, blackspot and powdery mildew in apples, and control of bacterial canker *Pseudomonas syringae* *pv.* *actinidiae* (Psa) in kiwifruit (**limited claim only**).



10L



9 418768 000001

FUNGICIDE/BACTERICIDE

Active Ingredient: Contains 5 g/litre Salicylic acid, 50 g/litre Benzalkonium chloride and 15 g/litre Propiconazole in the form of a Soluble Concentrate

Batch:

DOM:



Registered to:
Omnia Primaxa Ltd. T/A Omnia
Freephone 0800 774 629
www.omnia.net.nz
PO Box 201041
Auckland Airport, Manukau 2150

20L



9 418768 000018

IN A TRANSPORT EMERGENCY: DIAL 111. In case of emergency, phone 0800 243 622 or 0800 CHEMCALL



READ THE LABEL CAREFULLY BEFORE USING SPOTLESS



Active Ingredient: Contains 5 g/litre Salicylic acid, 50 g/litre Benzalkonium chloride and 15 g/litre Propiconazole in the form of a Soluble Concentrate

HSNO CLASS: 6.1D, 6.3A, 6.5A, 6.5B, 6.9B, 8.3A, 9.1A, 9.3C

TOXICITY:

TOXIC: MAY CAUSE RESPIRATORY SENSITISATION.
DANGER: THIS PRODUCT IS CORROSIVE AND MAY CAUSE EYE DAMAGE.

HARMFUL:

- MAY BE HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN.
- MAY CAUSE SKIN IRRITATION.
- MAY CAUSE SENSITISATION FROM PROLONGED SKIN CONTACT.
- MAY CAUSE ORGAN DAMAGE FROM REPEATED ORAL EXPOSURE AT HIGH DOSES.

ECOTOXICITY:

- VERY TOXIC TO FISH WITH LONG LASTING EFFECTS. AVOID CONTAMINATION OF ANY WATER SUPPLY WITH PRODUCT OR EMPTY CONTAINERS.
- HARMFUL TO TERRESTRIAL VERTEBRATES.

PRECAUTIONS:

- CORROSIVE - KEEP OUT OF REACH OF CHILDREN.
- When handling, wear eye protection, waterproof gloves, boots and overalls.
- Avoid contact with skin, eyes and inhalation of spray mist.
- Remove protective clothing and wash hands and face thoroughly before meals and after work.
- Do not eat, drink or smoke while using.

STORAGE:

- Store in the original container in a locked, dry, cool, well ventilated area out of direct sunlight, away from feed, seeds and foodstuffs. Avoid storage at temperatures below 5°C and above 35°C.
- Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

SHELF LIFE:

- When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DISPOSAL:

- Triple rinse empty container and add rinsate to the spray tank.
- Burn empty container in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush and bury empty container in suitable landfill. Do NOT use the container for any other purpose.
- Avoid contamination of any water supply with chemical or empty container.

SPILLAGE:

- Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste container. Wash area with water and absorb with further inert material. Dispose of waste safely (such as a suitable landfill).

FIRST AID:

If swallowed, DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.
If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately in running water for several minutes. If inhaled move the patient to fresh air immediately.

GENERAL DIRECTIONS:

Before using SPOTLESS carefully read the warning statement, precautions, instructions, first aid instructions and directions for use.

For optimum results SPOTLESS should be used as a preventative spray application.

RECORD KEEPING:

Written record of each application or discharge must be kept if 3 litres or more of the substance is applied in 24 hours in a place where the substance is likely to enter air or water and leave the place.

APPROVED HANDLER: Spotless must be under the personal control of an approved handler when the substance is:

- Applied in a wide dispersive manner, OR
- Used by a commercial contractor.

MIXING:

Shake the SPOTLESS container well before use. Add the required quantity of SPOTLESS to the spray tank partly filled with water and maintain gentle agitation while filling and spraying.

APPLICATION:

The dosage rates are recommended for high volume application to the point of run off. For concentrate spraying increase rates accordingly. The required dosage rate per hectare on trees should be determined by calculating the required water volume per hectare based on the tree row volume method.

COMPATIBILITY:

Can be tanked mixed with most commonly used fungicides and insecticides but do not mix with strong alkaline materials.

WITHHOLDING PERIODS:

Apples - Do not apply after the end of December or 60 days before harvest.
Kiwifruit - Do not apply after fruitset.
Olives - Apply post harvest and before fruitset the following season.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

It is an offence to use this product on animals

RESISTANCE MANAGEMENT:

Fungicides: Spotless contains propiconazole which is a DMI fungicide. Resistance to this fungicide may develop from excessive use. To minimise this risk use strictly in accordance with the label instructions and resistance management strategies that exist for any fungus listed on the label. Avoid using this fungicide exclusively all season.

Bactericides: Spotless contains benzalkonium chloride which has bactericidal properties. Resistance to this bactericide may develop from excessive use. To minimise this risk use strictly in accordance with the label instructions. Avoid using this bactericide exclusively all season.

DIRECTIONS FOR USE

Crop	Disease	Rate	Remarks
Apples	Blackspot and powdery mildew	500 ml / 100 litres of water	Apply Spotless at 10-14 day intervals from 75% petal fall to early December, ensuring a maximum of no more than four applications. A 7 day interval is recommended when conditions are conducive to severe black spot infection and during periods of rapid tree growth.
Kiwifruit	<i>Sclerotinia</i> fruit rot	350 ml / 100 litres of water (minimum of 3.5 litres in 1000 litres of water / ha)	Apply during flowering only. Do not apply after fruitset.
	Limited claim for the control of bacterial canker <i>Pseudomonas syringae</i> pv. <i>actinidiae</i> (Psa)	400 ml / 100 litres of water (minimum of 4 litres in 1000 litres of water / ha)	Apply at 10-14 day intervals from greentip to the end of flowering. Do not apply after fruitset. The efficacy of this product for control of Psa is currently supported by limited laboratory and greenhouse data and is therefore not fully proven at this stage. Warning: While crop-safety issues are not expected field data has not been provided to support the earlier timing between greentip and flowering.
Olives	Leafspot (Peacock spot)	350 - 500 ml / 100 litres of water	Commence spraying immediately after harvest and repeat at monthly intervals. Do not apply after fruitset the following season. Use the higher rate where a severe infection has occurred and apply at 14-21 day intervals until the infection is under control.

CONDITIONS OF SALE: The use of SPOTLESS being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Omnia accept no responsibility whatsoever resulting from the use of this product.

Registered pursuant to the ACVM Act 1997, No P7937. See <http://eatsafe.nzfsa.govt.nz/web/public/acvm-register> for registration conditions. Approved pursuant to the HSNO Act 1996, approval code HSR007857. See www.ermanz.govt.nz/search-databases/Pages/controls-search.aspx for controls applying to the substance.