

Molyphos

250g/L molybdenum 110g/L phosphorus as a mineral concentrate

Crop	Application rate per hectare	When to apply	Water rate litres/ hectare
Brassicas	300 to 500 mls	Soil applied prior to emergence; and	200 to 400
	100 to 400 mls	One of more foliar applications 8 to 12 days apart on young leaves.	200 to 400
Curcubits	100 to 250 mls	10 to 14 days after germination and at 4 to 5 leaf stage.	100 to 200
Lettuce	200 to 500 mls	Apply to young growth every 8 to 12 days as needed.	200 to 300
Lucerne	100 to 400 mls	Once or twice per season as necessary during strong growth after cutting.	100 to 200
Peas & Beans	300 to 500 mls	Soil applied prior to emergence; and	100 to 200
	200 to 500 mls	One application on young growth with a repeat in 8 to 12 days if necessary.	100 to 200
Pasture	50 to 200 mls	Apply while pasture is growing in Spring or Autumn.	100 to 200

Application rates vary. Refer to your Omnia representative for recommendations.

Caution: Excessive Molybdenum is toxic to plants and animals. Withhold stock from treated pasture for two weeks. Apply to no more than one fifth of total grazed area per year.

Conditions of sale: The use of this product being beyond the control of the suppliers, no warranty expressed or implied is given by Omnia regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and Omnia accept no responsibility whatsoever resulting from the use of this product.

Batch:



www.omnia.net.nz Freephone 0800 774 629
PO Box 201041 Auckland Airport, 2150

